

Selective Endoscopic Discectomy™ : Twelve Years Experience

Anthony T. Yeung, M.D.

Arizona Institute for Minimally Invasive Spine Care

Voluntary Associate Clinical Professor

Department of Orthopedics

University of California San Diego School of Medicine

Chapter Outline

- I. Introduction to SED**
- II. The Role of Evocative Chromo-Discography**
- III. The Role of Electro-thermal Therapy**
- IV. The Role of Laser**
- V. Adjunctive Therapy with Chymopapain**
- VI. Neuromonitoring**
- VII. Foraminal Epidurography and Therapeutic Injections**
- VIII. Present Applications and Future Trends**

Introduction

More than Twelve years have passed (1991) since I learned and adopted Arthroscopic Microdiscectomy from pioneer endoscopic spine surgeon Parviz Kambin, who, along with Sadahisa Hijikata, first established the technique for percutaneous nucleotomy in the early 1970's. A cadaver dissection of the traditional posterior anatomy of the lumbar spine compared with the foraminal anatomy clearly illustrates the feasibility and advantages of the foraminal

approach to the lumbar disc. **(fig 1a, b)** Kambin's Arthroscopic microdiscectomy technique evolved gradually to allow for more dorsal placement of the cannula to effect posterolateral fragmentectomy, resection of the posterior annulus for stenosis, and made possible the removal of extruded and sequestered herniated discs.[1-4] **(Fig 2. The dome)** Hijikata also recently reviewed his 12 years experience with endoscopic discectomy [5] In 1996, a new design of the operating spine endoscope adding multi-channel irrigation and complementary instrumentation **(Fig. 3 YESS scope)** [6] allowed further development of endoscopic spine surgery to include the treatment of annular tears causing discogenic back pain. [7, 8] Advanced techniques of foraminoplasty for central and lateral recess stenosis followed. [9-12] The third-generation system design change to the YESS system gave me more flexibility to maneuver the endoscope and improved ability to probe spinal anatomy in a conscious patient. [13] Continued evolution of this technique afforded me the ability to better evaluate the pathologic process causing the patient's discogenic back pain. Conditions previously not considered surgical, such as annular tears were evaluated and managed successfully through the endoscope. Synovial facet cysts, inflammatory membranes containing neo-neurogenesis, osteophytosis impinging on sensitive nerves, anomalous furcal nerves, autonomic nerves, and conditions irritating the dorsal root ganglion of the exiting nerve were identified as sources of discogenic pain and sciatica. From 1991 to 2004, I have treated over 2300 patients with discogenic pain, degenerative conditions of the lumbar spine, and the whole spectrum of disc herniations including extruded and sequestered fragments.[14, 15] The success rate in the first 500 patients was 432 (86%) of 500 good/excellent results according to the modified MacNab criteria. [11]A subsequent retrospective study of 219 consecutive patients with radiculopathy secondary to large intracanal non-contained lumbar disc

herniations demonstrated a satisfactory outcome in 204 (93.1%) based on modified MacNab criteria but the rate was even higher (94.8%) when patients were asked to respond to a study patient--based outcome questionnaire.[16] My 12 year's experience evolving from Kambin's arthroscopic microdiscectomy, into the treatment of discogenic back pain and sciatica by selective endoscopic discectomy, is reviewed .

The Role of Evocative Chromo-discography

At Kambin's 1991 course, Professor Hans Joerg Leu described the use of indigocarmine dye to stain and label the nucleus pulposus. In order to maintain the ability to recognize structural anatomy, it was necessary to dilute the dye to a 10% solution to effect differential tissue staining that did not overwhelm the nucleus pulposus with stain. I adopted this adjunctive technique initially to help with a visualized nucleotomy. [17] With differential staining, it was easier to recognize nucleus pulposus from annulus and from facet capsule. The epidural space with its epidural vessels and fat was simple to recognize. When pain was reproduced by discography, the clinical improvement of the patient's back pain correlated well with concordant pain reproduction. The use of discography also helped predict whether the herniation was extruded or contained, and the nuclear material was clearly stained for easier endoscopic extraction. [18] **(fig 4 discogram)**. I trademarked evocative chromodiscography™ as an integral part of spinal endoscopy. The process of removing the indigocarmine dye labeled nucleus was trademarked selective endoscopic discectomy™ to describe the technique as the focus of this chapter.[15] **(fig 5 nucleus pulposus stained with indigocarmine dye)** The literature on discography is currently considered controversial only because of the high inter-observer variability by discographers in reporting the patient's subjective pain as well as the ailing

patient's ability to give a clear response, especially if the pain response is altered by the use of analgesics or sedation during the procedure. Although much of the literature that contributes to the controversy of discography points out the pitfalls of depending on discography, the majority of the literature supports its use by clinicians who know how to use it. The surgeon who is accomplished in endoscopic spine surgery prefers to do the discography himself in order to decrease the interobserver variability in interpreting the patient's response. When we compare our own assessment of the patient's pain response with the report of another discographer, there can be some variability in diagnosis and interpretation. This variability may result in unpredictable treatment results. The incidence of "false positive" discograms, however, can be significantly decreased in the hands of an experienced endoscopic surgeon. "False positive" discography should really be false interpretation of positive discography results. The experienced endoscopic surgeon learns to correlate the patient's response to the discogram pattern of the painful disc that is being treated. There is good correlation of discograms with different types of annular tears and disc herniations. The surgical result can then be predicted on the basis of the visualized condition. For example, the discogram can be used to predict the presence of a collagenized disc fragment versus a soft herniation, the extrusion of a disc fragment as a non-contained herniation, or the presence of the type, grade, and location of a painful versus non-painful annular tear. Discography is used by the surgeon as a means of further identifying concordant discogenic pain in clinical situations where the patient's clinical presentation is matched with MRI findings. Controversy in the literature has arisen because of the lack of a good spectrum of therapeutic surgical treatments once the pain is confirmed, and because of the plethora of articles pointing out the pitfalls of false interpretation. [19, 20]

The Role of Electrothermal Therapy

Prior to my adoption of Arthroscopic microdiscectomy, I was using the KTP laser for laser disc decompression. When I combined the two techniques for nucleotomy, the laser provided hemostasis and better visualization. [21-24] I observed that patients with disc protrusions but with predominant back pain who were not candidates for traditional transcanal surgery found relief of their back pain with KTP assisted Arthroscopic microdiscectomy. The staining of the disc provided a chromophore that enhanced the efficacy of the KTP laser. [25] In 1993, I tested a unipolar electrode by Ellman International and introduced to me by Peter Morrison, M.D. Later, an electrode made by Smith and Nephew was used with the Ellman unit and the new working channel endoscope. In a study of my first 100 patients by Farouq Al-Hamdan , a spine fellow under Alex Hadjipavlou, it was documented in a retrospective 2 - 4.5 year follow-up study that the use of the KTP laser as an adjunct to Arthroscopic Micro-discectomy relieved back pain as well as leg pain in 65% of the patients. Overall good / excellent results by MacNab criteria was 89% for sciatica. The KTP laser was initially used to provide hemostasis and better visualization, but its side effect of laser thermal annuloplasty prompted an IRB study using a temperature controlled flexible probe by Oratec in lieu of the laser.

In 1998, an IRB approved study commenced at St Luke's Medical Center to evaluate the efficacy of electrothermal treatment in the process of arthroscopic microdiscectomy for herniated discs. This study, sponsored by Oratec, using a temperature controlled flexible probe, provided a better tool used to contract annular defects caused by the disc herniation and ablating granulation tissue in annular tears. The patient's response to this application of electro-thermal energy provided information that electrothermal treatment of the disc was effective in decreasing

discogenic back pain, but the fluctuations in temperature control caused me to switch to a bipolar flexible probe. [26, 27] Oratec investigated the IDET catheter in the same time period. I now use a bipolar flexible probe by Ellman designed specifically for thermal annuloplasty that has proven to be as effective as the Oratec probe, but of a more established and accepted use of electrothermal energy in spine surgery.[27] (**Fig 6 Ellman Triggerflex Probe**)

Rauschnig's cryosections of normal and patho-anatomy has more recently demonstrated inflammation in and around the sensitive dorsal root ganglion, and identified granulation tissue in annular tears. [28, 29](**fig 7 Annular tear**) (**fig 8 Exiting nerve root**) It is also well known that although a spinal structure is capable of pain, spinal pathology on imaging studies does not always correlate with the debilitating pain that is resistant to conservative management.[30] What may be very painful in one person may be well tolerated or painless in another. Evocative discography was shown to be helpful in identifying the disc as a pain generator in axial back pain and sciatica, [18, 31-37] and electrothermal treatment of the disc is demonstrated to be effective in decreasing discogenic back pain.

The Role of Laser

Laser technology also evolved to become more user friendly. The first Laser to be FDA approved was the KTP laser, a laser that was in the Blue/Green wave length spectrum that was effective for soft tissue ablation, but the bright light limited its use when direct visualization was desired.[22] The Holmium:yag laser was effective for the ablation of soft tissue as well as bone. Current designs now include a side firing irrigated probe and a straight fiber that can be delivered through a curved guide that will angle the laser beam up to 45 degrees. The laser has opened the door for the removal of osteophytes and lateral stenosis that cause neuropathic pain in patients

who have no other surgical options. [15] (**Figures 9-10**) demonstrate the use of laser in foraminoplasty.

Adjunctive therapy with Chymopapain

Chymopapain is the only minimally invasive technique that has been validated with two large double blind studies and numerous cohort studies that found it effective for the treatment of contained disc herniations. I have used chymopapain to assist the extraction of large non-contained disc herniations that extrude past the outer annular fibers, with a good /excellent result by MacNab criteria 10% higher than when no chymopapain is used. [38] If the herniation height is greater than the base on MRI, it is likely the herniated nucleus is collared by the annulus, making it more difficult to remove from within the disc. Chymopapain treated nucleus pulposus is soft and slippery, making mechanical removal easier. If chymopapain extravasates along the course of the contrast agent used for discography, it will theoretically denature and treat the extruded fragment to make it less inflammetogenic. The extruded fragment is also exposed to phagocytosis and eventual absorption if exposed to the epidural vasculature. In over 500 surgical cases of chymopapain assisted selective endoscopic discectomy, I have never had any complications from the use of chymopapain. Because of the absence of complications when used in conjunction with endoscopic disc removal, I now do not find it necessary to routinely test for antibodies to chymopapain with the Chymofast test unless the patient requests it. (**fig 11-13**)

Neuromonitoring

In one hundred consecutive cases, neuromonitoring was studied to determine whether complications sensory/motor complications could be reduced. I specifically studied whether neuromonitoring by continuous EMG or SSEP would affect the incidence of dysesthesia or help

predict decompression of the nerve.[39] While an increase conduction velocity was identified when a mechanically compromised nerve was decompressed, It was concluded that neuromonitoring was not any more effective than monitoring the patient for intra-operative pain when a dilute solution of lidocaine (.5%) was used. There was no difference in the dysesthesia or complication rate of the one hundred cases versus a matched number of patients who had no neuromonitoring. While neuromonitoring may give the novice surgeon a greater sense of security early in his endoscopic practice, I have not found it useful enough to consider it to be required for standard of care. Neuromonitoring may make surgeons feel more comfortable to warn him of nerve irritation from the procedure when he first practices endoscopic spine surgery, but after analyzing the result of the prospective study of 100 consecutive patients, it has not been shown to be any more useful than patient feedback reporting pain during the procedure for avoiding complications.

Foraminography and Therapeutic Injections

The efficacy of Endoscopic lumbar disc surgery can be enhanced by the surgeon performing his own discography and foraminal injections. Foraminal epidurography, and foraminal therapeutic injections are correlated with identification of patho-anatomy in the lumbar spine. (**fig 14-17 epidurograms**) Information gleaned from these injection procedures are used by the surgeon to better select patients for surgical interventional techniques that the surgeon has incorporated into his endoscopic surgical practice. Patients with disc protrusions, annular tears, and foraminal stenosis may get temporary relief with the therapeutic injection, but if the response is short lived, additional information gathered by performing the epiduralgram will help guide the surgeon when he has to consider the technical feasibility of using the same or similar portal for the

insertion of the operating cannulas. By performing epidurograms, the surgeon can get additional information of the anatomy of the foramen, the outline of the traversing and exiting nerves, and the therapeutic response afforded by the epidural injection.[37] The injection at L5-S1 will also help the surgeon with pre-operative planning if surgery is eventually required.

Present Applications and Future Trends

Indications

Any pathologic lesion that is accessible, visible, treatable, or requires endoscopic confirmation through the foramen may ultimately become an indication for diagnostic and therapeutic endoscopy. Patient selection for pain and radiculopathy from disc herniation is similar to selection criteria for traditional spine procedures. Endoscopic surgical indications, however, may be dictated by the limitations of the endoscopic procedure itself with respect to the patient's anatomy or the surgeon's skill and experience with endoscopic spine surgery. At L5-S1, anatomic restrictions may cause the surgeon to opt for the posterior transcanal approach (**Fig. 18 a,b, Narrow pelvis or horizontal disc**). For herniations from T10 to L4, the foraminal approach provides excellent access to the disc and epidural space. As the experience of the surgeon increases, previous contraindications become relative, depending partly on the surgeon's ability to endoscopically visualize, probe, and access the pathologic lesion. Restrictions are dictated only by anatomic considerations in accessing the patient's spinal pathology and the rationale for the endoscopic procedure itself. As the surgeon's experience increases, former contraindications became relative, depending on the surgeon's experience, and his or her ability to address the spinal condition to be treated.

The three zones within reach of the spine endoscope transforaminally are illustrated in **Fig. 19**

Inclusion Criteria

Discogenic pain as determined by evocative discography implicates the disc as a pain generator. Symptomatic disc herniation is the obvious indication, limited only by the accessibility of endoscopic instruments to the herniated fragment. The ideal lesion for endoscopic discectomy is a far lateral, extraforaminal disc herniation. Traditional approaches to far lateral disc herniations are more difficult, requiring a paramedian incision through very vascular tissue. The exiting nerve and the dorsal root ganglion is at risk for neuropraxia in both approaches.[28, 29] Although a traditional spine surgeon can access the lateral zone of the disc with a paramedian incision, it is easier to access the extraforaminal zone through the foramen. A typical foraminal view of nucleus pulposus extruded past the posterior annulus is shown in **(Fig 20)**. Through this approach to the disc, endoscopic excisional biopsy and disc space debridement are also ideal for surgically debriding infectious discitis **(Fig. 21)**. Currently treated with immobilization and parenteral antibiotics, discitis is much more effectively treated when augmented by endoscopic excisional biopsy and debridement. The surgeon will not have to be overly concerned about creating dead space for the inflamed or infected disc material to spread into the dead space created by a posterior approach. The clinical results are dramatic, and tissue biopsy is more accurate than needle aspiration in identifying the cause of discitis. Even sterile discitis will benefit from intradiscal debridement and irrigation.

Foraminal stenosis in selected patients will respond to foraminoplasty by endoscopic techniques. Lateral recess stenosis is one cause of failed back surgery syndrome (FBSS) that can absolutely be diagnosed and treated by foraminal decompression. [40] The patho-anatomic finding may be osteophytosis tethering the exiting nerve at the superior vertebral endplate and/or

stenosis and lack of fat around the exiting nerve. **(fig. 22)** Although trephines, rasps, and burrs can be used, the Ho:Yag side firing laser has made it feasible as a visually controlled soft tissue and bone ablation device. The cannula chosen for this task has an open side channel that will protect the exiting nerve while the laser is used to dissect the tethering osteophyte and scar tissue from the nerve. Endoscopic foraminoplasty has not been shown to cause increased instability even in spondylolisthesis. When mild degenerative spondylolisthesis is present, the disc bulge can be successfully treated by selective discectomy and thermal annuloplasty when there is a sciatic component to the patient's complaint. The technique is most useful for lateral recess stenosis, a condition that is responsible for atypical leg pain rather than true intermittent claudication of central spinal stenosis. In central spinal stenosis, when there is concomitant posterior disc protrusion, decompression of the spinal canal can be effectively accomplished by resecting the bulging annulus in a collapsed disc, thus lowering the floor of the foramen. In isthmic spondylolisthesis, when there is more leg than back pain, this is usually due to impingement on the exiting nerve by the pseudoarthrosis at the pars defect. The goal is then to decompress the compromised exiting nerve by elevating the dome formed by the inferior facet and lamina without further destabilizing the spinal segment.

Exclusion Criteria

Except for pregnancy, there are no absolute exclusion criteria but only relative contraindications depending on the skill and experience of the surgeon. Spinal endoscopy and spinal probing can be used for diagnostic purposes in very difficult or confusing clinical situations. Therefore, if endoscopy is helpful for diagnostic purposes, exclusion criteria may depend mainly on the accessibility of the spinal pathology and the endoscopic skills of the surgeon. The risks and

benefits of the procedure must be weighed against the need to use this fluoroscopically guided procedure under local anesthesia or sedation.

Future Considerations

The spine scope will eventually be used for any condition where visual inspection of the foramen is desired. I have used spinal endoscopy for the following reasons: (1) to inspect a spinal nerve that is suspected of being irritated by orthopedic hardware adjacent to the pedicle; (2) to remove suspected recurrent or residual disc herniations that do not show up on imaging studies; (3) to decompress the lateral recess by foraminoplasty; (4) to remove osteophytes and facet cysts that cause unrelenting sciatica; and (5) to locate painful lateral annular tears or small disc herniations not evident on physical examination or MRI. In Single and multi-level discogenic pain, where the patient has no viable options, endoscopic discectomy and thermal annuloplasty has been successful for treating chronic lumbar discogenic pain. A minority of patients may continue to have significant back pain, and a few may feel worse, but in the context of a progressive degenerative condition, the results are encouraging, and will give most patients relief while awaiting the development of newer procedures such as nucleus replacement, total disc replacement and minimally invasive stabilization procedures of the posterior spinal column.

Current Imaging Methods

In my experience, imaging studies are only about 70% accurate and specific for predicting pain. Conditions such as lateral annular tears, rim tears, endplate separation, small subligamentous disc herniations, intranuclear herniations, anomalous nerves, and miscellaneous discogenic conditions are cumulatively missed approximately 30% of the time. These conditions can be diagnosed by means of spinal endoscopy. Tears that are in the lateral and anterior aspect of the disc are

routinely missed on MRI studies. Very small disc herniations that protrude past the outer fibers of the annulus are also missed because the fragment may be flattened against the posterior longitudinal ligament or nerve, appearing on the MRI as a thickened or bulging annulus, but really contains a subligamentous herniation. When the nerve root is “swollen” or enlarged, the MRI is not always capable of distinguishing a swollen nerve from a conjoined nerve or a nerve with an adherent fragment of disc. When the disc tissue is in direct contact with the nerve, the nerve can be irritated and a painful inflammatory membrane forms. Even an epidural venous plexus that is inflamed can contribute to back pain and sciatica. Anomalous nerve branches known as furcal nerves are never seen on MRI but can be visualized with spinal endoscopy of the foramen.


When an inflammatory membrane is present, the patient’s pain pattern can be confusing. Diagnostic spinal endoscopy has confirmed “nondermatomal” pain in scores of patients with proximal thigh, buttock, and groin pain at levels distal to the root origin of the anatomic area.

Technique


Accessing the foramen is simplified and standardized by drawing coordinates on the patient’s skin to determine the optimal skin window and annular window for positioning the surgical instruments toward the center of the disc (**Fig. 23**). Reference points are the anatomic center of the disc, the superior facet of the inferior vertebra, and the skin window. The needle trajectory must also be in a line of inclination between the endplates of the adjacent vertebrae. Adjustments in the trajectory will be made to accommodate individual anatomic considerations and the pathology to be accessed. Once the optimal trajectory is established, the cannulas are inserted to allow for endoscopic surgery under direct visualization.

Endoscopic spine surgery has a very high learning curve but is within the grasp of every endoscopic surgeon with proper training. As with any new procedure, the complication rate is higher during the learning curve, and it may vary according to the skill and experience of each surgeon. The endoscopic technique, because of its approach, may pose additional risk for iatrogenic injury, but it is possibly safer than traditional surgery because the patient is awake and able to provide immediate input to the surgeon when pain is generated. The surgeon's ability to perform the surgery without causing the patient undue pain will self-select for surgeons who can master the technique to the extent that the surgeon prefers endoscopic surgery to traditional surgery for the same condition. For most disc herniations and discogenic pain, experienced endoscopic spine surgeons will opt for the endoscopic approach as the treatment of choice for their patients.

The future of endoscopic spine surgery is extremely bright. There will soon be an explosion of new imaging systems, endoscopes, and endoscopic instruments. Refined techniques and image-guided systems may help diminish the learning curve. Coupled with advancements in tissue regeneration and enhancement of tissue healing, and the trend toward tissue healing instead of removal, regeneration over healing, and arthroplasty instead of fusion, the spine surgeon may no longer have to consider spine surgery as paradoxical. As a treatment modality, it will no longer be considered a last resort in a desperate patient. There will be a paradigm shift in the way we view and approach patients with back pain, especially when endoscopic spine surgery is further validated with outcome studies and becomes routinely available.


LEGENDS**Fig. 1 a**

Anatomy of the posterior port provides easier access to the posterior disc and spinal canal at L5-S1 (blue hubbed needle), but with planning, most contained disc herniations can be removed posterolaterally.

Fig 1 b


Anatomy of the posterolateral foraminal port from L2-S1. Only in the L5-S1 disc space is access to the spinal canal restricted because of the pelvis and the relatively wide facet (gray hubbed needle in the L5-S1 disc) High lumbar disc herniations from L1 to L3 are easier to reach endoscopically through the posterolateral foraminal portal. L4-5 provides ample room for either approach. Note the furcal nerve branches entering the psoas muscle.

Fig. 2. The dome.



Spinal structures in the foramen accessible to visualization and surgical intervention and probing via the posterolateral approach includes the annulus, disc, pedicle, facet, and epidural space. This approach also visualizes neuro-compression in the the “hidden zone” of the lateral recess, a common cause of failed back surgery syndrome. (courtesy of Hal Matthews, M.D.).

Fig 3 Yeung Spine Scope System



The Yeung Endoscopic Spine System discoscope and partial instrument set. The spinal endoscope is designed with multichannel irrigation and a cannula system that allows access to targeted areas while protecting sensitive nerves. (From Yeung AT, Yeung, CA. Advances in endoscopic disc and spine surgery: Foraminal approach. In Szabó I, Coburg AJ, Strange PS, et al., eds. Surgical Technology International. XI. San Francisco: Universal Medical Press, 2003, p 257.)

Fig 4. Discogram



This Intraoperative discogram in a patient with a papacentral HNP by Mri not only confirmed the patient's concordant back pain and sciatica, but the leakage of contrast to the traversing and exiting nerves alerted the surgeon to look for extruded disc fragments and grade V annular tears, especially in the lateral zone.

Fig. 5 Nucleus pulposus stained with Indigocarmine dye



The Indigocarmine dye stains the nucleus pulposus light blue, helping the endoscopic surgeon target the stained nucleus for extraction. Here, the herniation has extruded past the unstained


annulus. Removal of the extruded herniation will expose the traversing nerve in the epidural space.

Fig. 6 Ellman Triggerflex probe




This bipolar flexible device has provided effective hemostasis, electrothermal shrinkage of disc tissue, and annular tears under visualized control. It offers better control of the energy source by its bipolar design.

Fig. 7. A, Annular tears.



This CT discogram outlines a foraminal HNP and far lateral annular tear that will irritate the dorsal root ganglion of the exiting nerve. Endoscopic visualization of the foramen may reveal the presence of an inflammatory membrane, extruded nucleus pulposus, and granulation tissue


Fig. 7B



Granulation tissue and inflammation surrounding a small disc fragment in the foramen.

Grade V annular tears open into the epidural space or psoas muscle, allowing the in growth of nerves and capillaries that create an inflammatory response, which, if next to a spinal nerve or the dorsal root ganglion, can cause pain out of portion to what may be anticipated from traditional imaging studies. Tears in this area will also result in groin pain by irritating the psoas muscle and genital-femoral nerve. If patients with annular tears obtain relief from foraminal epidural blocks, more lasting relief of 2 or more years has been documented with selective endoscopic discectomy and thermal annuloplasty.

Fig 7 C



Bipolar radiofrequency treatment of annular tears under direct visualization. Interpositional disc material should be removed from the annular layers to effectively treat the annular tear.

Fig 7 D



Preoperative endoscopic view of a grade V annular tear demonstrating granulation tissue in the tear.

Fig 7E



Postoperative view of the annulus after thermal modulation with the Ellman Bipolar Triggerflex Probe.


Fig 8 The Exiting Nerve



The exiting nerve is in the “hidden extraforaminal zone” that is irritated by far lateral annular tears and disc herniations that escape detection by MRI and transcanal surgical exploration. It contains the Dorsal Root Ganglion, that, when sensitized by the inflammatory by-products of a degenerative disc, is responsible for the “nondermatomal “ distribution in patients with chronic


sciatica. The presence of fat around the exiting nerve is a more sensitive indication of lateral recess stenosis than findings on MRI or CT myelogram.

Fig. 9 The Role of laser in Foraminoplasty




Side firing laser (Trimedine) or end firing laser fibers (Lisa) directed by a flexible cannula
Are used for precise foraminoplasty of the lateral recess.

Fig. 10 a



Post-op Endoscopic view of foraminoplasty of the superior articular facet at L5-S1

Fig 10b



Post-op view of decompressed exiting nerve after foraminoplasty

Fig. 11-13 Chymopapain assisted SED








Figure 11 a, b

Pre-op MRI of an extruded fragment in a horizontal disc at L5-S1 that is anatomically difficult to reach. By using chymopapain, the results of endoscopic removal has been demonstrated to be improved by 10% overall, by reducing the rate of residual HNP or recurrent HNP.


Figure 12



Extruded fragment labeled by indigocarmine and pre-treated by chymopapain allowed for easier manual extraction. The chymopapain loosened the fragment made it easier to remove, aided by suction on the working channel of the endoscope.

Figure 13

Chymopapain treated nucleus pulposus. Note that the differential staining of the nuclear material extracted in the above patient. Unstained collagenized disc was extracted from the epidural space where the indigocarmine dye did not reach. By visualizing the decompressed foramen, successful relief of leg pain was immediate post –op.

Fig 14 Foraminal epidurograms

Foramino-epidurography: a new technique for foraminal needle placement from the far lateral skin portal mimicking the surgical access to the epidural space allows the surgeon to produce an epidurogram that complements the MRI by outlining the position of the traversing and exiting nerves in the foramen. This information provides the surgeon additional information pre-operatively and serves as a “practice run” for surgery.

Fig 15a ,b. Anatomic limitations At L5-S1




The narrow pelvis in this patient limited access to the disc, but it was still possible to get a needle into the disc for discography. Depending on the type of disc protrusion, the endoscopic surgeon will have a better idea about the prognosis of the surgical procedure contemplated. If this patient had an extruded disc herniation, it would be better to opt for the trans-canal approach.

Figure 15 b Horizontal L5-S1 Disc




The pelvis becomes even a greater obstacle, at L5-S1 if the disc is horizontal, making it much more difficult to get into the epidural space. Removal of the lateral facet could overcome this obstacle. Good pre-op planning is enhanced by foramino-epidurography.

Fig 16 Three zones on the foramen accessible by the endoscope




Zones two and three are not usually visualized by surgeons using the trans-canal approach unless they are experienced in the paramedical approach to the lateral recess.

Fig 17 Foraminal view of HNP




The indigocarmine stained disc tissue has extruded past the posterior longitudinal ligament in this foraminal HNP at L4-5

Fig. 18. Discitis



Intradiscal view of discitis after debridement. Usual findings of inflammatory disc material and loose endplate cartilage are readily removed from the disc space. Pain relief is immediate, and abundant tissue is available for laboratory analysis.


Fig 19 Dissecting exiting nerve



In lateral recess stenosis, the scarred exiting nerve is being released with a bare laser fiber. This picture demonstrates lateral recess stenosis as the most common cause of Failed Back Surgery Syndrome.

Fig 20 YESS Technique

DR. YEUNG'S AMD INSTRUMENTATION TRAJECTORY PROTOCOL



Determination of optimal instrument path: Yeung instrumentation trajectory protocol.

Interoperative C-arm fluoroscopic imaging allows registration of internal structures with surface skin markings. **A**, Posteroanterior fluoroscopic exposure enables topographic location of spinal column midline and transverse planes of target discs. Intersections of drawn lines mark posteroanterior disc centers. **B**, Lateral fluoroscopic exposure enables topographic location of the lateral disc center and allows visualization of the plane of inclination for each disc. **C**, The inclination plane of each target disc is drawn on the skin from the lateral disc center to the posterior skin surface. **D**, The distance between the lateral disc center and the posterior skin

surface plane is measured along each disc inclination line. **E** and **F**, This distance is then measured from the midline along the respective transverse plane line for each disc. At the end of this measure a line parallel to midline is drawn to intersect each disc inclination line. This intersection marks the skin entry point of “skin window” for each target disc. Needle insertion at this point toward the target disc at an angle 25 to 30 degrees to the surface skin plane will determine the path of all subsequent instrumentation.

Bibliography

1. Kambin, P., *Arthroscopic microdiscectomy*. Spine J, 2003. **3**(3 Suppl): p. 60S-64S.
2. Kambin, P., et al., *Transforaminal arthroscopic decompression of lateral recess stenosis*. J Neurosurg, 1996. **84**(3): p. 462-7.
3. Kambin, P., et al., *Arthroscopic microdiscectomy and selective fragmentectomy*. Clin Orthop, 1998(347): p. 150-67.
4. Kambin, P. and M.H. Savitz, *Arthroscopic microdiscectomy: an alternative to open disc surgery*. Mt Sinai J Med, 2000. **67**(4): p. 283-7.
5. Hijikata, S., *Percutaneous nucleotomy. A new concept technique and 12 years' experience*. Clin Orthop, 1989(238): p. 9-23.
6. Yeung, A.T. *Spinal endoscopy with a multichannel, continuous irrigation, discoscope with integrated inflow and outflow ports (poster presentation)*. in *North American Spine Society Annual Meeting*. 1997. New York, New York.
7. Yeung, A.T. *Arthroscopic Electro-thermal Surgery for Discogenic Low Back Pain: a Preliminary Report*. in *International Intradiscal Therapy Society Annual Meeting*. 1998. San Antonio, Texas.
8. Yeung, A.T. *Classification and Electro-Thermal Treatment of Annular Tears*. in *American Back Society Annual Meeting December 12*. 1998. Las Vegas, Nevada.
9. Yeung, A.T. *Minimal Access Surgery for Degenerative Conditions of the Lumbar Spine*. in *3rd Congress Meeting of the Chinese Speaking Orthopedic Society and Annual Congress of the Chinese Orthopedic Association*. 1999. Taipei, Taiwan, October 13-15.
10. Yeung, A.T., *Minimally Invasive Disc Surgery with the Yeung Endoscopic Spine System (YESS)*. Surg Technol Int, 2000. **VIII**: p. 267-277.
11. Yeung, A.T., *The evolution of percutaneous spinal endoscopy and discectomy: state of the art*. Mt Sinai J Med, 2000. **67**(4): p. 327-32.

12. Yeung, A.T., *Selective Discectomy with the Yeung Endoscopic Spine System*, in *The Practice of Minimally Invasive Spinal Technique*, M.H. Savitz, J. Chiu, and A.T. Yeung, Editors. 2000, AAMISMS Education LLC. p. 115-122.
13. Yeung, A.T. *Patho-anatomy of Discogenic Pain*. in *Minimally Invasive Spine Update*. 1999. Disney Magic Cruise Nov 5-8.
14. Yeung, A.T. and P.M. Tsou, *Posterolateral endoscopic excision for lumbar disc herniation: Surgical technique, outcome, and complications in 307 consecutive cases*. *Spine*, 2002. **27**(7): p. 722-31.
15. Yeung, A.T. and C.A. Yeung, *Advances in endoscopic disc and spine surgery: foraminal approach*. *Surg Technol Int*, 2003. **11**: p. 253-61.
16. Tsou, P.M. and A.T. Yeung, *Transforaminal endoscopic decompression for radiculopathy secondary to intracanal noncontained lumbar disc herniations: outcome and technique*. *Spine J*, 2002. **2**(1): p. 41-8.
17. Yeung, A.T. *The value of an intra-operative discogram: Its role in Arthroscopic microdiscectomy*. in *International Intradiscal Therapy Society Annual Meeting and San Francisco Spine Institute Course*. 1995.
18. Yeung, A.T., *The role of provocative discography in endoscopic disc surgery*, in *The Practice of Minimally Invasive Spinal Technique*, M.H. Savitz, J. Chiu, and A.T. Yeung, Editors. 2000, AAMISMS Education LLC. p. 231-236.
19. Carragee, E.J., et al., *The rates of false-positive lumbar discography in select patients without low back symptoms*. *Spine*, 2000. **25**(11): p. 1373-80; discussion 1381.
20. Carragee, E.J., et al., *False-positive findings on lumbar discography. Reliability of subjective concordance assessment during provocative disc injection*. *Spine*, 1999. **24**(23): p. 2542-7.
21. Yeung, A.T. *Enhancement of KTP/532 LDD and AMD Procedures with a vital Dye*. in *AAOS Annual Meeting Scientific Exhibit*. 1992. San Francisco Feb 18-23.
22. Yeung, A.T., *Considerations for Use of the KTP Laser for Disc Decompression and Ablation*, in *Spine: State of the Art Reviews*. 1993.
23. Yeung, A.T. *Enhancement of KTP/532 LDD and AMD Procedures with a Vital Dye*. in *International Intradiscal Therapy Society March 10-14*. 1993. Phoenix, Arizona.
24. Yeung, A.T., *Enhancement of KTP/532 Laser Disc Decompression and Arthroscopic Microdiscectomy with a Vital Dye*. 1994, AAOS Audio Video Program, AAOS 61st Annual Meeting: New Orleans, Louisiana.
25. Yeung, A.T. *Laser as an adjunct to Arthroscopic Microdiscectomy*. in *North American Spine Society Annual Meeting*. 1997. New York, New York.
26. Yeung, A.T. and S.A. Gore, *Evolving methodology in treating discogenic back pain by Selective Endoscopic Discectomy (SED)*. *Journal of Minimally Invasive Spinal Technique*, 2001. **1**: p. 8-16.
27. Yeung, A.T., et al., *Intradiscal thermal therapy for discogenic low back pain*, in *The practice of minimally invasive spinal technique*, M.H. Savitz, J. Chiu, and A.T. Yeung, Editors. 2000.

28. Yeung, A.T. *Macro-and Micro-anatomy of Degenerative Conditions of the Lumbar Spine (Best Paper Presentation Award)*. in *International Intradiscal Therapy Society 16th Annual Meeting*. 2003. Chicago, Illinois April 2-5.
29. Yeung, A.T. *Rauschnig's Anatomy for Minimally Invasive Spine Surgery*. in *Spine Across the Sea 2003 July 27-31*. 2003.
30. Modic, M.T., et al., *Imaging of degenerative disk disease*. Radiology, 1988. **168**(1): p. 177-86.
31. Aprill, C. and N. Bogduk, *High-intensity zone: a diagnostic sign of painful lumbar disc on magnetic resonance imaging*. Br J Radiol, 1992. **65**(773): p. 361-9.
32. Aprill, C.N., *Diagnostic Disc Injection*, in *The Adult Spine: Principles and Practice*, F. J.W., Editor. 1996, Lippincott-Raven Publishers: Philadelphia. p. 539-562.
33. Bini, W., et al., *The role of provocative discography in minimally invasive selective endoscopic discectomy*. Neurocirugia (Astur), 2002. **13**(1): p. 27-31; discussion 32.
34. Guyer, R.D. and D.D. Ohnmeiss, *Lumbar discography*. Spine J, 2003. **3**(3 Suppl): p. 11S-27S.
35. Guyer, R.D. and D.D. Ohnmeiss, *Lumbar discography. Position statement from the North American Spine Society Diagnostic and Therapeutic Committee*. Spine, 1995. **20**(18): p. 2048-59.
36. Yeung, A.T. *Evocative Lumbar Discography: Treatment of Multi-level cases*. in *1st World Congress American Academy of Minimally Invasive Spinal Medicine and Surgery*. 2000. Las Vegas, Nevada Dec 7-10.
37. Yeung, A.T. *Discography, Foraminal Epidurography and Therapeutic Foraminal Injections: Its Role in Endoscopic Spine Surgery*. in *International 22nd course for Percutaneous Endoscopic Spinal Surgery and Complementary Techniques*. 2004. Zurich, Switzerland January 20-30.
38. Yeung, A.T. *Intra-operative Chemoneucleolysis as an adjunct to Arthroscopic Microdiscectomy*. in *International Intradiscal Therapy Society Annual Meeting*. 1997. Naples, Florida.
39. Yeung, A.T., J. Porter, and C. Merican. *SEP as a sensory integrity check in selective endoscopic discectomy using the Yeung endoscopic spine system*. in *2nd World Congress American Academy of Minimally Invasive Spinal Medicine and Surgery*. 2001. Las Vegas, Nevada December 2001.
40. Yeung, A.T., *Failed Back Surgery Syndrome*, in *The Practice of Minimally Invasive Spinal Technique*, M.H. Savitz, J. Chiu, and A.T. Yeung, Editors. 2000, AAMISMS Education LLC. p. 293-296.

SUGGESTED READING

1. Aprill C, Boduk N. High-intensity zone: A diagnostic sign of painful lumbar disc on magnetic resonance imaging. Br J Radiol 65:361-369, 1992.

2. Freemont AJ. Nerve ingrowth into diseased intervertebral disc in chronic back pain. *Lancet* 350:178-181, 1997.
3. Hermantin FU, Peters T, Quararo L, Kambin P. A prospective, randomized study comparing the results of open discectomy with those of video-assisted arthroscopic microdiscectomy. *J Bone Joint Surg Am* 81:958-965, 1999.
4. Javid MJ, Nordby EJ, Ford LT, et al. Safety and efficacy of chymopapain in herniated nucleus pulposus with sciatica. *JAMA* 249:2489-2494, 1983.
5. Jensen MC, Brant-Zawadzki, MN, Obuchowski N, et al. Magnetic resonance imaging of the lumbar spine in people without back pain. *N Engl J Med* 331:69-73, 1994.
6. Kambin P, Gennarelli T, Hermantin F. Minimally invasive techniques in spinal surgery: Current practice. *Neurosurg Focus* 4:1, 1998.
7. Kambin P, O'Brien E, Zhou L, Schaffer JL. Arthroscopic microdiscectomy and selective fragmentectomy. *Clin Orthop Rel Res* 347:150-167, 1998.
8. Kambin P. Posterolateral percutaneous lumbar discectomy and decompression: Arthroscopic microdiscectomy. In Kambin P, ed. *Arthroscopic Microdiscectomy: Minimal Intervention in Spinal Surgery*. Baltimore: Urban & Schwarzenberg, 1991, pp 67-100.
9. Kuslich SD. Microsurgical lumbar nerve root decompression utilizing progressive local anesthesia. In Williams W, McCullouch J, Young P, eds. *Microsurgery of the Lumbar Spine*. Rockville, MD: Aspen Publishing, 1990.
10. Matthews HH, Long BH. *Principles and Practice of Spine Surgery*. Philadelphia: Mosby, 2003, pp 288-294.

11. Osman SG, Nibu K, Panjabi MM, Marsolais EB, Chaudhary R. Transforaminal and posterior decompression of the lumbar spine. A comparative study of stability and intervertebral foramen area. *Spine* 22:1690-1695, 1997.
12. Osti OL, Vernon-Roberts B, Moore R, Fraser RD. Annular tears and disc degeneration in the lumbar spine. A post mortem. *J Bone Joint Surg Br* 74:678-682, 1992.
13. Saal JA, Saal JS, Ashley J. Targeted intradiscal thermal therapy: Preliminary feasibility studies. ISSLS Conference, June 2-3, 1997.
14. Sachs B, Vanharanta H, Spivey MA, et al. Dallas discogram description: A new classification of CT/discography in low back pain disorders. *Spine* 12:287-294, 1987.
15. Savitz MH, Chiu JC, Yeung AT. *The Practice of Minimally Invasive Spinal Technique*, 1st ed. Richmond, VA: AAMISMS Education, LLC: 1-368, 2000.
16. Schaffer J, Kambin P. Percutaneous posterolateral lumbar discectomy and compression with 6.9 millimeter cannula: Analysis of operative failures and complications. *J Bone Joint Surg Am* 73:882-831, 1991.
17. Schellhas KP, Pollei SR, Gundry CR, Heithoff KB. Lumbar disc high-intensity zone: Correlation of magnetic resonance imaging and discography. *Spine* 21:79-86, 1996.
18. Tsou PM, Yeung AT. Transforaminal endoscopic decompression for radiculopathy secondary to intracanal noncontained lumbar herniations. Out come and technique. *Spine* 27:41-48, 2002.
19. Tsou PM, Yeung CA, Yeung AT, "Selective Endoscopic Discectomy™ and Thermal Annuloplasty for Chronic Lumbar Discogenic Pain: a minimal access visualized intradiscal procedure" *The Spine Journal* (Accepted 1-04 in press July/August 04)

20. Yeung AT, Gore SA. Evolving methodology in the treatment of discogenic back pain by selective endoscopic discectomy. *J Minimally Invasive Spinal Technique* 1:8-14, 2001.
21. Yeung AT. Minimally invasive disc surgery with the Yeung Endoscopic Spine System (Y.E.S.S.). *Surgical Technology International VIII*, June, 1999.
22. Yeung AT, Yeung CA. Advances in endoscopic disc and spine surgery: Foraminal approach. *Surgical Technology International XI*, June 253-61, 2003
23. Yeung AT. The evolution of percutaneous spinal endoscopy and discectomy: State of the Art. *Mt Sinai J Med* 67:327-332, 2000.
24. Yeung AT, Tsou PM. Posterolateral endoscopic excision for lumbar disc herniation. Surgical technique, outcome, and complications in 307 consecutive cases. *Spine* 27:722-731, 2002.