

coflex® Device and Procedure

For decades, LSS patients' surgical options were limited to either decompression or decompression with spinal fusion. In 2012, the FDA approved the **coflex®** spinal implant, which is a small, U-shaped titanium device that provides spinal stability without the mobility loss associated with spinal fusion.

How is the **coflex®** device implanted?

After the surgical decompression, which removes pressure on the impinged nerves, your surgeon will insert the **coflex®** implant through the same incision. The implant is then positioned onto the lamina, which is the strongest bone in the back of your spine.

The unique design of the **coflex®** implant maintains stability in the spine while preserving more natural movement at the affected area.

How do patients with **coflex®** compare to patients with spinal fusion?

A 2013 published study comparing the **coflex®** device with spinal fusion found that the **coflex®** device had "advantages in perioperative outcomes," and that "equivalent or superior 2-year clinical outcomes were seen with **coflex®**." The article concluded that "**coflex®** is a safe, efficacious, and viable alternative to spinal fusion in the treatment of spinal stenosis with low back pain."¹⁻³

What are the benefits of **coflex®** vs. decompression with pedicle screws?

In comparative research vs. spinal fusion, the data demonstrates that **coflex®** patients do better, faster.¹

Patients receiving the **coflex®** implant experienced faster relief of their symptoms, reported higher satisfaction rates with their outcomes, and required less post-operation pain medication than spinal fusion patients.²


Decompression with the **coflex®** device is faster than decompression with pedicle screws, causes less blood loss and requires a shorter hospital stay.³

Is the **coflex®** device the right choice for me?


To be a candidate for treatment for decompression with the **coflex®** device, you must be skeletally mature and must have moderate-to-severe spinal stenosis in your lower back. Symptoms of LSS include difficulty walking a long way, such as half a mile, and having pain in your lower back while standing that subsides when you bend forward.

In order to receive the **coflex®** implant, you must have been treated by a doctor for at least six months with non-surgical treatments.

Traditional Option: Spinal Fusion



New Option: **coflex®** Interlaminar Stabilization™



¹⁻³ Davis RJ, Errico TJ, Bae H, Auerbach JD (2013): Decompression and Coflex® Interlaminar Stabilization Compared With Decompression and Instrumented Spinal Fusion for Spinal Stenosis and Low-Grade Degenerative Spondylolisthesis. Spine 2013; 38: 1529-1539.